CONTENTS

	Page
Flavor of Roquefort Cheese. JAMES N. CURRIE	1
Corn-Leaf Blotch Miner. W. J. PHILLIPS	15
Coloration of the Seed Coat of Cowpeas. ALBERT MANN	33
Experiments with Apple Leaf-Spot Fungi. JOHN W. ROBERTS	57
Longevity of Pycnospores of the Chestnut-Blight Fungus in Soil. F. D. HEALD and M. W. GARDNER	67
Behavior of Cup Current Meters Under Conditions not Covered by Standard Ratings. FRED C. SCOBEY	77
Eremocitrus, A New Genus of Hardy, Drouth-Resistant Citrous Fruits from Australia. WALTER T. SWINGLE	85
Relation of Bacterial Transformations of Soil Nitrogen to Nutrition of Citrous Plants. KARL F. KELLERMAN and R. C. WRIGHT.	101
Aroma of Hops: A Study of the Volatile Oil with Relation to the Geographical Sources of the Hops. FRANK RABAK	115
Lesser Bud-Moth. E. W. SCOTT and J. H. PAINE	161
Two New Wood-Destroying Fungi. JAMES R. WEIR	163
Bacteria Concerned in the Production of the Characteristic Flavor in Cheese of the Cheddar Type. ALICE C. EVANS, E. G. HASTINGS, and E. B. HART	167
Relation of the Action of Certain Bacteria to the Ripening of Cheese of the Cheddar Type. E. B. HART, E. G. HASTINGS, E. M. FLINT, and ALICE C. EVANS	193
Citrus-Root Nematode. N. A. COBB	217
Pellicularia Koleroga on Coffee in Porto Rico. G. L. FAWCETT	231
Feeding Habits of the Boll Weevil on Plants Other than Cotton. B. R. COAD	235
Identity of Peridermium Fusiforme with Peridermium Cerebrum. GEORGE G. HEDGCOCK and W. H. LONG	247
Identification of Species of Fusarium Occurring on the Sweet Potato, Ipomoea Batatas. H. W. WOLLENWEBER	251
Mutation in Egyptian Cotton. THOMAS H. KEARNEY	287
Influence of the Host on the Morphological Characters of Puccinia Ellisiana and Puccinia Andropogonis. W. H. LONG	303
Ability of Streptococci to Survive Pasteurization. S. HENRY AYERS and WILLIAM T. JOHNSON, Jr	321

Page

Fruit-Rot, Leaf-Spot, and Stem-Blight of the Eggplant Caused by Phomopsis Vexans. L. L. HARTER 331
Head Smut of Sorghum and Maize. ALDEN A. POTTER 339
Oxidases in Healthy and in Curly-Dwarf Potatoes. H. H. BUNZEL 373
Birds as Carriers of the Chestnut-Blight Fungus. F. D. HEALD and R. A. STUDHALTER 405
Density of Wood Substance and Porosity of Wood. FREDERICK DUNLAP 423
Composition of Roquefort-Cheese Fat. JAMES N. CURRIE 429
A New Sarcophagid Parasite of Grasshoppers. E. O. G. KELLY .. 435
Papaya Fruit Fly. FREDERICK KNAB and W. W. YOTHERS 447
Index 455

ERRATA

Page 108, line 22, "between 0.05 and 0.0100 per cent" should read "between 0.05 and 0.100 per cent."

Page 217, heading, "The Citrus-Root Nematode Tylenchus Semipenetrans" should read "The Citrus-Root Nematode Tylenchulus Semipenetrans."

ILLUSTRATIONS

PLATES

Corn-Leaf Blotch Miner

Page

PLATE I. Fig. 1.—Large corn leaf, showing blotch mines. Fig. 2.—Young corn leaf from breeding cage containing six larvæ. Fig. 3.—Shows how mines occasionally fill with water. Fig. 4.—Millet leaf showing feeding punctures. Fig. 5.—Large corn leaf with three larvæ in the same tunnel. Fig. 6.—Section of leaf shown in figure 5 slightly enlarged. Fig. 7.—Straight mine ending in a slight blotch in a large leaf..... 32

PLATE II. Fig. 1.—Rearing shelter used for *Agromyza parvicornis* and other insects at La Fayette, Ind. Fig. 2.—Rearing cage used for *Agromyza parvicornis*..... 32

PLATE III. Fig. 1.—*Agromyza parvicornis*, the corn-leaf blotch miner: *A*, Dorsal view of adult; *B*, antenna of female; *C*, head of male; *D*, hypopygium of male; *E*, ovipositor of female. Fig. 2.—*Pleurotropis utahensis*, a parasite of *Agromyza parvicornis:* Adult..... 32

PLATE IV. Fig. 1.—*Closterocerus tricinctus*, a parasite of *Agromyza parvicornis:* *A*, Dorsal view of adult; *B*, side view of antenna. Fig. 2.—*Opius diastatae*, a parasite of *Agromyza parvicornis:* Adult. Fig. 3.—*Opius succineus*, a parasite of *Agromyza parvicornis:* Adult..... 32

PLATE V. Fig. 1.—*Derostenus punctiventris*, a parasite of *Agromyza parvicornis:* Adult. Fig. 2.—*Opius utahensis*, a parasite of *Agromyza parvicornis:* Adult..... 32

Coloration of the Seed Coat of Cowpeas

PLATE VI. Fig. 1.—Transverse section of the seed coat of a cowpea, similar to that shown in text figure 1, but showing the relative thickness of three layers, as on the seed. The cells are not expanded with chloral hydrate. *cu*, Cuticle; *a*, palisade layer; *b*, middle or hour-glass layer; *c*, basal-color layer. Fig. 2.—Seeds of cowpeas, showing some of the variations in the style of marking of the seed coat..... 56

Experiments with Apple Leaf-Spot Fungi

PLATE VII. Fig. 1.—Apple leaves from Tennessee, showing typical spots of the frog-eye disease. Fig. 2.—Spores of *Alternaria mali*, which is capable of enlarging dead spots on apple leaves. Fig. 3.—York Imperial apple leaves, showing spots enlarged by *Alternaria mali*..... 66

Eremocitrus, a New Genus of Hardy, Drouth-Resistant Citrous Fruits from Australia

PLATE VIII. *Eremocitrus glauca* (Lindl.) Swing. Grown from seed from the Condamine River near Chinchilla, Queensland, Australia. Fig. 1.—Desert kumquat grafted on the Australian finger lime (*Citrus australasica*). Fig. 2.—Desert kumquat grafted on the wood-apple (*Feronia elephantum*).. 100

TWO NEW WOOD-DESTROYING FUNGI

Page

PLATE IX. Fig. 1.—*Fomes putearius* from the wood of *Larix occidentalis*, showing a typical pileate fruiting structure; also a section through the resupinate form. Fig. 2.—*Larix occidentalis*, showing decay of the wood by *Fomes putearius*. Fig. 3.—*Pinus ponderosa*, showing disintegration of the wood by *Fomes putearius*. Fig. 4.—*Pseudotsuga taxifolia*, showing the typical decay caused by *Fomes putearius*. Fig. 5.—*Picea engelmanni*, showing the different stages of the decay caused by *Fomes putearius* 166

PLATE X. Fig. 6.—*Trametes setosus* on *Pinus monticola;* resupinate form. Fig. 7.—*Trametes setosus* on *Pinus monticola;* sessile pileate form. Fig. 8.—*Trametes setosus:* common form on *Thuya plicata*. Fig. 9.—*Trametes setosus* on *Larix occidentalis*. Fig. 10.—*Thuya plicata*, showing decay of the wood caused by *Trametes setosus*. Fig. 11.—*Trametes setosus:* Transverse section of the pores, showing the numerous setæ 166

IDENTITY OF PERIDERMIUM FUSIFORME WITH PERIDERMIUM CEREBRUM

PLATE XI. Fig. 1.—A globoid swelling formed by *Peridermium cerebrum* on *Pinus echinata*, showing the cerebroid arrangement of the æcia. Fig. 2.—A fusiform swelling caused by *Peridermium fusiforme* on *Pinus taeda*, showing a similar cerebroid arrangement of the æcia. Fig. 3.—A leaf of *Quercus rubra* bearing the telia of *Cronartium cerebrum* from an inoculation with æciospores of *Peridermium cerebrum* on *Pinus virginiana*. Fig. 4.—A leaf of *Quercus phellos* bearing the uredinia of *Cronartium cerebrum* from an inoculation with the æciospores of *Peridermium fusiforme* 250

IDENTIFICATION OF SPECIES OF FUSARIUM OCCURRING ON THE SWEET POTATO, IPOMOEA BATATAS

PLATE XII. Fig. A.—*Fusarium batatatis*, n. sp.: Mycelium stage with blue sclerotial plectenchymata on steamed potato tuber. Fig. B.—*Fusarium batatatis*, n. sp.: Ochreous- to salmon-colored pionnotes on the same medium, but from a transfer of a single sickle-shaped conidium. Fig. C.—*Fusarium batatatis*, n. sp.: Wine-red acid color modification of the fungus on steamed rice, turning blue with alkali 286

PLATE XIII. Figs. A–G.—*Hypomyces ipomoeae* (Hals.) Wollenw. Fig. H.—*Fusarium incarnatum* (Rob.) Desm. Fig. J.—*Hypomyces cancri* (Rutg.), n. comb. Fig. A.—*Hypomyces ipomoeae:* Ascospores. 1, Ellipsoidal shape; 2, form of a double paraboloid; 3, overripe, slightly swollen stage; 4, separation of the two cells in overripe stage. Fig. B.—*Hypomyces ipomoeae:* Asci with a paraphysis. Fig. C.—*Hypomyces ipomoeae:* Perithecium. Fig. D.—*Hypomyces ipomoeae:* Chlamydospores. 1 and 3, lateral; 2, intercalated and terminal; 4, intercalated within a conidium (conidio-chlamydospore). Fig. E.—*Hypomyces ipomoeae:* Perithecium formed by spiral coiling of a lateral hypha. Fig. F.—*Hypomyces ipomoeae:* False conidial heads produced at the end (1) of conidiophores by spores suspended in drops of water. The conidiophores sprang from an old conidium which was separated into two parts, one of which was dead (*a*). Fig. G.—*Hypomyces ipomoeae:* Normal conidia. 1, More curved at the ends than the other spores; 2–7, tri- to quinque-septate spores. Fig. H.—*Fusarium incarnatum* (Rob.) Desm.: Conidia, *a*, short, *b*, slender; 1–3, normal 5-septate conidia; 4, lanceolate; 5, exceptionally large; *c*, pedicellate base without heel. Fig. J.—*Hypomyces cancri* (Rutg.), n. comb.: Mature conidia, the first one (1) being especially large 286

Page

PLATE XIV. *Gibberella Saubinetii* (Mont.) Sacc. Figs. A–C.—*Gibberella Saubinetii:* Perithecia grown in pure culture. Fig. A.—On stem of *Vicia faba*, with two ostiola surrounded by a collar of large peridial cells. FIG. B.—On wheat grains showing the ascus ball after one-half of the peridium had been lifted by a longitudinal section. Fig. C.—On Irish potato stem, without a distinct collar. Fig. D.—*Gibberella Saubinetii:* Two asci with a paraphysis. Fig. E.—*Gibberella Saubinetii:* Ascospores, 1, with slightly swollen cells, overripe; 2, normal shape; 3, dried condition. Figs. F–G.—*Gibberella Saubinetii:* Normal conidia grown on Irish potato stem. Fig. F.—12 days old. Fig. G.—6 days old, culture watered more; therefore the conidia are broader than in figure F. Fig. H.—*Gibberella Saubinetii:* Abnormally multiseptate conidium about to germinate. These conidia are frequently to be found with swollen cells in young, and sometimes old cultures on the parenchyma of potato tubers. Fig. J.—*Gibberella Saubinetii:* Plectenchymatic parts of a stroma, formed by closely interwoven chains of swollen cells which have a thick membrane and brown to red contents with many vacuoles. Fig. K.—*Gibberella Saubinetii:* A number of conidia formed in young cultures on the moist surface of potato cylinders. Septation (1, 3) and shape (2) rarely normal; 4, mother conidium broken into two halves (a, b), both of which have developed some small conidiophores from the cells torn asunder 286

PLATE XV. Fig. A.—*Hypomyces ipomoeae* (Hals.) Wollenw.: Sweet-potato strain. 1, 2, Grown in pure culture on cotton stem; 3, on maple stem; 4, on wheat straw; 5 and 6, on potato cylinder. Fig. B–C.—*Hypomyces cancri* (Rutg.), n. comb.: The hemp strain found with perithecia on a dead taproot of hemp at the soil level. Fig. B.—Grown on steamed corn kernels. Fig. C.—1, Perithecia from the original field material; 2, 3, Grown in pure culture on cotton stem. Figs. D–G.—*Gibberella Saubinetii* (Mont.) Sacc.: The mycelium stage. Fig. D.—Grown on stem of Robinia, which was more or less reduced when single ascospores of these perithecia were transferred to fresh moist steamed stems of plants. Fig. E.—Grown on wheat straw. Figs. F and G.—Grown on cotton stem (on almost dry wheat straw the stroma could be reduced still more and completely disappeared in some of the later cultures on this medium) 286

PLATE XVI. *Fusarium* spp. on sweet potato with and without known perfect stage, grown on sterilized vegetables. Figs. A–E.—*Fusarium batatatis*, n. sp. Fig. A.—Microconidia. Fig. B.—1, Intercalated and terminal chlamydospores; 2, In young stage; 3, Branch from sclerotial plectenchymata; therefore no true chlamydospores; 4, Chlamydospores formed from the content of conidial cells (conidio-chlamydospores); 5, Mature chlamydospores. Fig. C.—Two conidia anastomosing, one of them producing microconidia. Fig. D.—Normal conidia from sporodochia. Fig. E.—Conidiophore from a sporodochium. Figs. F–P.—Characteristic conidia of different species of Fusarium. Fig. F.—*Fusarium hyperoxysporum*, n. sp. Fig. G.—*Fusarium acuminatum* Ell. and Ev. Fig. H.—*Hypomyces ipomoeae* (Hals.) Wollenw. Fig. J.—*Fusarium culmorum* (W. G. Sm.) Sacc. Fig. K.—*Fusarium radicicola*, n. sp. Fig. L.—*Fusarium incarnatum* (Rob.) Sacc. Fig. M.—*Fusarium caudatum*, n. sp. Fig. N.—*Fusarium orthoceras*, var. *triseptatum*, n. var. Fig. O.—*Gibberella Saubinetii* (Mont.) Sacc. Fig. P.—*Fusarium caudatum*, var. *volutum*, n. var. 286

MUTATION IN EGYPTIAN COTTON

Page

PLATE XVII. Field of Egyptian cotton (Gila variety) in blossom at the Cooperative Testing and Demonstration Gardens, Sacaton, Ariz., on July 15, 1913 302

PLATE XVIII. Fig. 1.—A plant of the Yuma variety of Egyptian cotton, photographed on July 15, 1913, showing the well-developed vegetative branches and the rather poorly developed lower fruiting branches. Fig. 2.—A plant of the Pima variety of Egyptian cotton, photographed on July 15, 1913, showing the absence of vegetative branches and the presence of well-developed fruiting branches at the low nodes on the axis. Fig. 3.—A plant of the Gila variety of Egyptian cotton, photographed on July 15, 1913, showing in comparison with the Yuma variety (Pl. XVIII, fig. 1) a smaller development of the vegetative branches 302

PLATE XIX. Leaves of the Yuma variety of Egyptian cotton taken from the main stem and showing the strong tendency in this variety to produce 5-lobed leaves 302

PLATE XX. Leaves of the Pima variety of Egyptian cotton taken from the main stem 302

PLATE XXI. Leaves of the Gila variety of Egyptian cotton taken from the main stem 302

PLATE XXII. Involucres of Egyptian cotton. Fig. 1.—Nubari variety. Fig. 2.—Yuma variety 302

PLATE XXIII. Involucres of Egyptian cotton. Fig. 1.—Pima variety. Fig. 2.—Gila variety 302

PLATE XXIV. Bolls of Egyptian cotton. Fig. 1.—Nubari variety. Fig. 2.—Yuma variety. Fig. 3.—Pima variety 302

PLATE XXV. Bolls of Egyptian cotton. Fig. 1.—Mit Afifi variety. Fig. 2.—Gila variety. Fig. 3.—Big-bolled strain of the Gila variety. Fig. 4.—Sakellaridis variety 302

FRUIT-ROT, LEAF-SPOT, AND STEM-BLIGHT OF THE EGGPLANT CAUSED BY PHOMOPSIS VEXANS

PLATE XXVI. "Damping off" of seedling eggplants which had been sprayed with organism No. 113 (*Phyllosticta hortorum*) 338

PLATE XXVII. Seedling eggplants from the pot shown in Plate XXVI, showing the presence of pycnidia and the effect of the fungus on the stem 338

PLATE XXVIII. An eggplant leaf sprayed with organism No. 104 (*Phoma solani*), showing the characteristic spots and pycnidia formed therein 338

PLATE XXIX. Fig. 1.—An eggplant fruit produced by a plant grown on the Potomac Flats and mummified as a result of spraying with organism No. 104 (*Phoma solani*). Fig. 2.—A photomicrograph of a cross section through pycnidia formed on the calyx of an eggplant 338

PLATE XXX. A section through the fruit of an eggplant which had been sprayed in a moist chamber with organism No. 113 (*Phyllosticta hortorum*) 338

HEAD SMUT OF SORGHUM AND MAIZE

Page

PLATE XXXI. Fig. 1.—Head smut in ear of maize. Fig. 2.—Head smut in tassel of maize 372

PLATE XXXII. Fig. 1.—The three American species of sorghum smut on Blackhull kafir: (a) *Sphacelotheca cruenta*, (b) *Sorosporium reilianum*, (c) *Sphacelotheca sorghi*. Fig. 2.—Head smut, *Sorosporium reilianum* (Kühn) McAlp., on "sumac" sorgo, San Antonio, Tex., October, 1913 372

PLATE XXXIII. Fig. 1.—Section through young sorus, showing hyphal aggregates preceding spore formation. Fig. 2.—Section through immature sorus. Note the fibrovascular bundle on the left, about which the spores, none of which were as yet quite mature, were developing in groups even in the earliest stages 372

PLATE XXXIV. Fig. 1.—Growth on carrot agar at various temperatures; cultures about 6 weeks old. Figs. 2 and 3.—Twenty-two days' growth on synthetic glucose agar (fig. 2) and on carrot agar (fig. 3). Fig. 4.—Chlamydospores formed in culture in peptonized maltose solution 372

PLATE XXXV. Fig. 1.—Smutted culms of Amber sorgo, showing the characteristic sterility of the main panicle. Fig. 2.—Proliferated head of Blackhull kaoliang, with one normal and one smutted head. Fig. 3.—Head-smut infection experiment. Control pot (see Table V, plat B, No. 8) on left; inoculated pot (Table V, plat B, No. 3) on right, showing three smutted plants 372

PLATE XXXVI. Panicular formation in apex of proliferated sorghum flower. Longitudinal section, showing presence of hyphæ of head smut 372

PLATE XXXVII. Longitudinal sections through the growing points of two of the buds indicated in text figures 1 and 2, showing hyphæ of the head smut. Fig. 1.—Bud 3 of culm B1. Fig. 2.—Bud 7 of culm N2 372

BIRDS AS CARRIERS OF THE CHESTNUT-BLIGHT FUNGUS

PLATE XXXVIII. Old blight canker on chestnut, showing the work of woodpeckers 422

PLATE XXXIX. Series of cultures 9 days old obtained in the test of bird No. 23, a brown creeper 422

A NEW SARCOPHAGID PARASITE OF GRASSHOPPERS

PLATE XL. Fig. 1.—*Sarcophaga kellyi:* Adult male fly, dorsal view. Fig. 2.—*Sarcophaga kellyi:* Head of male, front view. Fig. 3.—*Sarcophaga kellyi:* Head of male, lateral view. Fig. 4.—*Sarcophaga kellyi:* Hypopygium of male, lateral view. Fig. 5.—*Sarcophaga kellyi:* Left half of main forceps of same, ventrolateral view. Fig. 6.—*Sarcophaga kellyi:* Lateral view of same. Fig. 7.—*Sarcophaga kellyi:* Head of female, front view 446

PAPAYA FRUIT FLY

PLATE XLI. Fig. 1.—*Toxotrypana curvicauda* Gerst.: Female. Fig. 2.—*Toxotrypana curvicauda* Gerst.: Male 454

PLATE XLII. Fig. 1.—*Toxotrypana curvicauda* Gerst.: Eggs. Fig. 2.—*Toxotrypana curvicauda* Gerst.: Larva. Fig. 3.—*Toxotrypana curvicauda* Gerst.: Puparium. Fig. 4.—Papaya fruit infested with larvæ of *Toxotrypana curvicauda* Gerst. 454

TEXT FIGURES

Corn-Leaf Blotch Miner

Page

FIG. 1. Map of the United States, showing distribution of *Agromyza parvicornis*, the corn-leaf blotch miner 17

2. Tip of corn leaf, showing work of *Agromyza parvicornis*, the corn-leaf blotch miner: *a*, Egg; *b*, newly hatched larva in mine; *c*, feeding punctures of adults 19

3. *Agromyza parvicornis*, the corn-leaf blotch miner: Full-grown larva; *a. sp.*, anterior spiracles; *p. sp.*, posterior spiracles 20

4. *Agromyza parvicornis*, the corn-leaf blotch miner: Mouth hooks of larva. 20

5. *Agromyza parvicornis*, the corn-leaf blotch miner: Puparium; *a*, dorsal view; *b*, ventral view; *c*, view of posterior extremity 21

6. *Pleurotropis utahensis*, a parasite of *Agromyza parvicornis:* Larva 29

Coloration of Seed Coat of Cowpeas

FIG. 1. Transverse section of the seed coat of a cowpea with the cells expanded with chloral hydrate so as to show the structure of the three layers: *cu*, Cuticle; *a*, palisade layer; *b*, middle or hour-glass layer; *c*, basal-color layer 35

2. Outer surface of the seed coat of a cowpea, showing the upper ends of the palisade cells 37

Behavior of Cup Current Meters under Conditions not Covered by Standard Ratings

FIG. 1. Standard-rating curve on logarithmic scale, as developed by U. S. Bureau of Standards, for new cup meter, with points developed by rating the same meter at Calexico, Cal., under conditions similar to those under which it was originally rated and under other conditions not met in the usual rating 78

2. Rating curves and individual points on logarithmic scale, developed in the rating of a new cup meter at the hydraulic laboratory of Cornell University by holding the meter nearer the surface of the water and nearer the sides of the channel than is done in the usual rating 79

3. Vertical velocity curves, showing error due to using standard rating for points near the surface and bottom, Orr ditch, Nevada 80

4. Curves on logarithmic scale, showing relation between rate of vertical movement of a cup meter and velocity of water as indicated by the revolutions of the turbine due to vertical movement in still water .. 81

Eremocitrus, a New Genus of Hardy, Drouth-Resistant Citrous Fruits from Australia

FIG. 1. *Eremocitrus glauca:* Twigs from a syntypic specimen collected by T. L. Mitchell, near Forestvale, Queensland, Australia, on October 17, 1846; from Gray Herbarium. *A*, Spiny twig showing a single flower; *B*, spineless sterile twig; natural size. *C*, tetramerous flower seen from the side 86

2. *Eremocitrus glauca:* Fruits and seed from Tambo, Queensland, Australia. *A*, large pyriform fruit, natural size; *B*, small oblate fruit, natural size; *C*, cross section of a 4-celled fruit, showing four seeds and numerous pulp vesicles, natural size; *D*, a single, short-stalked pulp vesicle; *E*, seed, showing rugose testa 87

Page

FIG. 3. *Eremocitrus glauca:* Seedling plants grown from seed from near Chinchilla, Queensland, Australia. *A*, young seedling with hypogeous cotyledons still inclosed in the seed coats, natural size; *B*, an older seedling with a very long taproot, natural size; *C*, a cataphyll..... 88

4. *Eremocitrus glauca:* Flower from which petals and stamens have fallen; showing pedicel, calyx (one sepal cut away), disk, ovary, style, and stigma; from a specimen collected by Robert Brown, Sept., 1802 (?), Upper Head, Broad Sound, Queensland; in National Herbarium, Washington, D. C..... 89

5. *Eremocitrus glauca:* Cross section of leaf from near Forestvale, Queensland, collected by T. L. Mitchell on October 17, 1846; in Gray Herbarium..... 93

6. *Eremocitrus glauca:* A hair from the leaf of a plant grown from seed from near Chinchilla, Queensland..... 94

7. *Eremocitrus glauca:* Cross section of the epidermal region of a 1-year-old twig from near Forestvale, Queensland, collected by T. L. Mitchell on October 17, 1846; in Gray Herbarium. *A*, thick-walled epidermis, two strata of palisade cells, and inner cortical parenchyma; *B*, a stomate at the bottom of a deep narrow pit..... 94

RELATION OF BACTERIAL TRANSFORMATIONS OF SOIL NITROGEN TO NUTRITION OF CITROUS PLANTS

FIG. 1. Diagram showing relative nitrogen-fixing power, nitrifying power, and salt content of soil samples from good and poor areas in an orange grove which was unproductive and markedly chlorotic throughout. The good areas were themselves very unproductive. Black columns show good areas; hatched columns indicate poor areas..... 102

2. Diagram showing relative nitrogen-fixing power, nitrifying power, and salt content of good and poor areas in an orange grove that produced a fair crop of marketable fruit. The poor areas were unproductive. Black columns show good areas; hatched columns indicate poor areas. 103

3. Diagram showing relative nitrogen-fixing power, nitrifying power, and salt content of soil samples from good and poor areas in an orange grove which showed decided symptoms of chlorosis associated with small yields of low-grade fruit. There was but a slight difference in the vigor of the orange trees growing upon the good and the poor areas. Black columns show good areas; hatched columns indicate poor areas..... 103

4. Diagram showing quantities of salts found in crust samples of soil from the same areas illustrated in figures 1, 2, and 3..... 104

5. Diagram showing effect upon nitrate nitrogen in samples of California soil of additions of 2 per cent of green vetch, green barley, and mature barley straw..... 108

6. Diagram showing effect upon total nitrogen, not including nitrate nitrogen, in samples of California soil of additions of 2 per cent of green vetch, green barley, and mature barley straw..... 110

7. Diagram showing effect of additions of 2 per cent of green vetch, green barley, and mature barley straw upon the nitrifying power of samples of California soil..... 110

AROMA OF HOPS: A STUDY OF THE VOLATILE OIL WITH RELATION TO THE GEOGRAPHICAL SOURCES OF THE HOPS

Page

FIG. 1. Relative ester content of various hop oils..... 127
2. Fractionation curves of hop oils from the crops of 1906..... 130
3. Fractionation curves of hop oils from the crops of 1907..... 131
4. Fractionation curves of hop oils from the crops of 1908..... 132
5. Fractionation curves of hop oils from the crops of 1909..... 133
6. Optical-rotation curves of hop oils from the crops of 1906..... 137
7. Optical-rotation curves of hop oils from the crops of 1907..... 138
8. Optical-rotation curves of hop oils from the crops of 1908..... 139
9. Optical-rotation curves of hop oils from the crops of 1909..... 140
10. Ester-number curves of hop oils from the crops of 1906..... 143
11. Ester-number curves of hop oils from the crops of 1907..... 144
12. Ester-number curves of hop oils from the crops of 1908..... 145
13. Ester-number curves of hop oils from the crops of 1909..... 146
14. Fractionation curves of original and saponified hop oils..... 153

CITRUS-ROOT NEMATODE

FIG. 1. *Tylenchulus semipenetrans:* Mature and half-grown females, with their head ends permanently embedded in the feeding root of a Citrus tree. This is a plant parasite similar in many ways to the notorious gallworm *Heterodera radicicola*. *a*, Tail end; *b*, vulva; *c*, excretory pore; *d*, *e*, egg in uterus..... 218

2. Healthy Citrus root magnified 5 diameters. Note the size and form of the healthy root endings. Compare with figure 3..... 218

3. Citrus root attacked by the parasitic nematode *Tylenchulus semipenetrans*, magnified 5 diameters. The parasites are shown at *a*. They are shown black, but in reality are yellowish or brownish..... 219

4. *Tylenchulus semipenetrans:* Lateral view of full-grown male. *a*, Lip region; *b*, spear; *c*, 3-bulbed base of spear; *d*, œsophageal lumen; *e*, median œsophageal bulb; *f*, nerve ring; *g*, cardiac œsophageal bulb; *h*, beginning of the intestine; *j*, large intestinal granule; *k*, small intestinal granule; *l*, anus; *m*, spicula; *n*, excretory pore; *o*, spermatocyte; *p*, vas deferens; *q*, spermatozoon..... 220

5. *Tylenchulus semipenetrans:* Female ready for fertilization. *a*, Lip region; *b*, guide of spear; *c*, shaft of spear; *d*, lumen of œsophagus; *e*, median œsophageal bulb; *f*, nerve ring; *g*, cardiac œsophageal bulb; *h*, beginning of intestine; *o'*, ovary; *n'*, duct of renette; *n*, excretory pore; *r*, vulva; *s*, wings of the cuticle; *l'*, deteriorated anus; *i*, terminus..... 221

6. *Tylenchulus semipenetrans:* Larva, soon after hatching from the egg. *a*, Lip region; *b*, spear; *c*, 3-bulbed base of spear; *d*, lumen of the œsophagus; *e*, median œsophageal bulb; *f*, nerve ring; *g*, posterior œsophageal bulb; *h*, beginning of the intestine; *j*, large intestinal granule; *k*, small intestinal granule; *n*, excretory pore; *i*, terminus.. 222

7. *Tylenchulus semipenetrans:* Young larva of male undergoing the first or second molt. *a*, Lip region; *a'*, lip region of sloughed skin; *b*, deteriorating spear; *b'*, spear of the sloughed skin; *d*, lumen of the œsophagus; *e*, median œsophageal bulb; *f*, nerve ring; *g*, cardiac œsophageal bulb; *h*, beginning of the intestine; *j*, large intestinal granule; *k*, small intestinal granule; *n*, excretory pore; *o*, immature internal sexual organ; *l*, developing male sexual opening; *i*, terminus 223

Page

FIG. 8. Two half-sections of Citrus root magnified 30 diameters: *a*, Healthy root; *b*, diseased root 224

9. *Tylenchulus semipenetrans:* Posterior portion of adult female. The portion shown is located outside of the Citrus root. *a*, Blind end of the coiled single ovary; *b*, anterior portion of ovary; *c*, posterior portion of ovary; *d*, renette cell; *e*, duct of the renette cell; *f*, excretory pore; *g*, vulva; *h*, terminus; *i*, uterus; *j*, cuticle; *k*, somatic granule; *l*, junction of ovary with uterus; *m*, posterior end of ovary with ovum ready to enter uterus 224

10. Pot containing Citrus seedling. *a*, Brood capsule; *b*, label of same attached to capsule by means of a thread which passes down through the soil 225

11. Longitudinal section of a glass brood capsule enlarged 10 diameters. *a*, Glass tube shown in longitudinal section; *b*, thread encircling capsule, shown in cross section; *c*, finest mesh linen cloth held over bottom of filter capsule by the encircling cord *b*; *d*, layer of filter paper, for convenience shown black; *e*, *f*, disks of filter paper; *g*, nematode egg embedded in Pasteurized orange-root detritus and soil granules 225

12. *Mononchus papillatus* Bastian: Rather immature female specimen which has been feeding upon *Tylenchulus semipenetrans*. *a*, Two of the innervated papillæ existing on one of the six mobile lips; *b*, one of the lips; *c*, dorsal pharyngeal tooth; *d*, one of the three longitudinal chitinous ribs of the pharynx; *e*, pharyngeal cavity; *f*, œsophagus; *g*, muscular layer of the body; *h*, cuticle; *i*, one of the cells of the intestine; *j*, *j*, *j*, oral spears of three ingested Tylenchuli; the spear in the intestine near the vulva is accompanied by an undigested portion of the lumen of the œsophagus of the Tylenchullus; *k*, nerve ring; *l*, blind end of the anterior ovary, which, being behind the intestine, shows less clearly than the posterior ovary; *m*, nucleus of one of the intestinal cells; *n*, vulva; *o*, anus; *p*, terminus; *q*, anal muscles; *r*, rectum; *s*, cardia; *t*, spicula of an ingested male Tylenchulus 226

13. *Tylenchulus semipenetrans:* Young female about to undergo its second molt. *a*, Lip region; *b*, spear; *b'*, spear which has been sloughed off; *d*, lumen of the œsophagus; *e*, median œsophageal bulb; *f*, nerve ring; *g*, cardiac œsophageal bulb; *h*, beginning of the intestine; *k*, larger intestinal granules; *n*, excretory pore; *n'*, renette cell; *o'*, blind end of single immature ovary; *r*, developing uterus; *l'*, remnant of rudimentary anus; *i*, terminus 227

PELLICULARIA KOLEROGA ON COFFEE IN PORTO RICO

FIG. 1. Early stage in development of group of hold-fast cells of *Pellicularia koleroga* 231

2. Appressoria of Venezuelan candelillo fungus from an old, fully developed specimen 232

3. Spores of a fungus accompanying *Pellicularia koleroga* 232

ABILITY OF STREPTOCOCCI TO SURVIVE PASTEURIZATION

Page

FIG. 1. Results of heating streptococci for 30 minutes at various temperatures. 323
2. Results of heating streptococci (classified according to source) for 30 minutes at various temperatures. 324
3. Results of heating streptococci (classified as typical and atypical) for 30 minutes at various temperatures. 327

FRUIT-ROT, LEAF-SPOT, AND STEM-BLIGHT OF THE EGGPLANT CAUSED BY PHOMOPSIS VEXANS

FIG. 1. Some microscopic characters of the fungus *Phomopsis vexans: a*, Pycnospores; *b*, conidiophores; *c*, stylospores. 336

HEAD SMUT OF SORGHUM AND MAIZE

FIG. 1. Diagrammatic representation of the occurrence of infection in the nodal buds or branches of several sorghum plants. 346
2. Diagram of Plate XXXVII, figure 1, showing the position of the hyphæ. 347
3. Curves summarizing for different years the percentages of infection in plantings of sorgo after all-hot-water treatments and in control plantings. 349
4. Curves summarizing for different years the percentages of infection: First, in plantings of Red Amber sorgo after modified hot-water treatments at all temperatures and of all durations, but after presoakings of various duration; and, second, in control plantings (not treated with hot water). 350
5. Curves summarizing for different years the percentages of infection in plantings of Red Amber sorgo after modified hot-water treatments at all temperatures and of all presoakings, but after treatments of various duration. 351
6. Curves summarizing for different years the percentages of infection in plantings of Red Amber sorgo after modified hot-water treatments of all durations of presoaking and treatment, but at various temperatures. 352
7. Curves summarizing for different years the percentages of infection: First, in plantings of sorgo after all seed treatments; and, second, in control plantings. 353

OXIDASES IN HEALTHY AND IN CURLY-DWARF POTATOES

FIG. 1. Rate of development of the aerial portion of potato plants and mean rate of development of all the single shoots in one hill. 378
2. Curve showing oxidation of leuco base of malachite green in the presence of the juice of green potato shoots. 379
3. Curve showing oxidation of aloin in the presence of the juice of green potato shoots. 379
4. Curve showing oxidation of tyrosin in the presence of the juice of green potato shoots. 379
5. Curve showing oxidation of phloridzin in the presence of the juice of green potato shoots. 381
6. Curve showing oxidation of leuco base of malachite green in the presence of the juice of potato foliage. 382
7. Curve showing oxidation of pyrocatechol in the presence of the juice of potato foliage. 382

Page

FIG. 8. Curve showing oxidation of phloridzin in the presence of the juice of potato foliage 384
9. Curve showing oxidation of o-cresol in the presence of the juice of potato foliage 384
10. Curve showing oxidation of m-cresol in the presence of the juice of potato foliage 385
11. Curve showing oxidation of p-cresol in the presence of the juice of potato foliage 386
12. Curve showing oxidation of m-toluidin in the presence of the juice of potato foliage 386
13. Curve showing oxidation of pyrogallol in the presence of the juice of potato tubers 391
14. Curve showing oxidation of α-naphthol in the presence of the juice of potato tubers 393
15. Curve showing oxidation of tyrosin in the presence of the juice of potato tubers 393
16. Curve showing oxidation of hydrochinone in the presence of the juice of potato tubers 394
17. Curve showing oxidation of guaiacol in the presence of the juice of potato tubers 395
18. Curve showing oxidation of o-cresol in the presence of the juice of potato tubers 395
19. Curve showing oxidation of p-cresol in the presence of the juice of potato tubers 396
20. Curve showing oxidation of p-toluidin in the presence of the juice of potato tubers 397
21. Curve showing the differences in rate and extent of growth between healthy and diseased plants 398

BIRDS AS CARRIERS OF THE CHESTNUT-BLIGHT FUNGUS

FIG. 1. Spores other than those of *Endothia parasitica* obtained by the microscopic examination of the centrifuged sediment from the test of bird No. 7, a downy woodpecker: *a*, Brown; *b*, dark brown; *c*, brown; *d*, cyanophyceous; *e*, brown; *f*, nearly black; *g*, smoky; *h*, brown; *i*, brown; *j*, brown; *k*, brown; *l*, cyanophyceous 418
2. Spores other than those of *Endothia parasitica* obtained by the microscopic examination of the centrifuged sediment from the test of bird No. 23, a brown creeper. Brown-spored forms appear to predominate: *a*, Hyaline; *b*, hyaline; *c*, dark brown; *d*, nearly black; *e*, hyaline; *f*, light brown; *g*, smoky; *h*, pale smoky; *i*, dark smoky; *j*, hyaline; *k*, pale smoky; *l*, dark smoky; *m*, smoky; *n*, hyaline; *o*, brown; *p*, hyaline; *q*, dark smoky; *r*, hyaline; *s*, very dark, almost black 419